

Initial singlet and triplet spin state contributions to $\vec{p}\vec{p} \rightarrow pp\pi^0$

P. Thörngren Engblom^{a1}, H.O. Meyer^a, J.T. Balewski^a, W.W. Daehnick^b, J. Doskow^a,
W. Haeberli^c, B. Lorentz^c, P.V. Pancella^d, R.E. Pollock^a, B. von Przewoski^a,
F. Rathmann^e, T. Rinckel^a, Swapan K. Saha^b, B. Schwartz^c, A. Wellinghausen^a, T. Wise^c

^aIndiana University Cyclotron Facility, Indiana University, Bloomington, IN 47405, U.S.A.

^bDepartment of Physics, University of Pittsburgh, Pittsburgh, PA 15260, U.S.A.

^cDepartment of Physics, University of Wisconsin-Madison, Madison, WI 53706, U.S.A.

^dDepartment of Physics, Western Michigan University, Kalamazoo, MI 49008, U.S.A.

^eFriedrich-Alexander Universität, Erwin-Rommel Str. 1, 91508 Erlangen, Germany.

The PINTEX² facility at the IUCF Cooler ring, dedicated to the study of spin dependence in nucleon-nucleon interactions, has been used to measure polarization observables of the reaction $\vec{p}\vec{p} \rightarrow pp\pi^0$ at beam energies between 325 and 400 MeV. The stored, polarized proton beam had spin projections both in the longitudinal and the transverse directions with respect to the beam momentum. We report here on the measurements of the relative transverse and longitudinal spin-dependent cross sections³, $\Delta\sigma_T/\sigma_{tot}$ and $\Delta\sigma_L/\sigma_{tot}$, and how from these observables the initial spin singlet and triplet cross sections are obtained. Considering angular momentum states less than or equal to one, the contribution of the Ps partial waves to the cross section can be extracted.

1. INTRODUCTION

For reactions with two incoming spin 1/2 particles one can define the initial state cross sections $^{2S+1}\sigma_{m_s}$ where S denotes the combined spin of the two particles and m_s its projection. The three possible initial spin state contributions $^1\sigma_0$, $^3\sigma_0$ and $^3\sigma_1$ are related to $\Delta\sigma_T$ and $\Delta\sigma_L$ by isospin, parity and angular momentum conservation

$$\frac{^1\sigma_0}{\sigma_{tot}} = 1 + \frac{\Delta\sigma_T}{\sigma_{tot}} + \frac{1}{2} \frac{\Delta\sigma_L}{\sigma_{tot}} \quad (1)$$

$$\frac{^3\sigma_0}{\sigma_{tot}} = 1 - \frac{\Delta\sigma_T}{\sigma_{tot}} + \frac{1}{2} \frac{\Delta\sigma_L}{\sigma_{tot}} \quad (2)$$

$$\frac{^3\sigma_1}{\sigma_{tot}} = 1 - \frac{1}{2} \frac{\Delta\sigma_L}{\sigma_{tot}} \quad (3)$$

¹Present address: Department of Radiation Sciences, Box 535, S-75121 Uppsala, Sweden.

Email: pia.thorngren@tsl.uu.se

²<http://www.iucf.indiana.edu/~pintex/>

³Defined as $\Delta\sigma_{T(L)} = [\sigma(\uparrow\downarrow) + \sigma(\downarrow\uparrow)] - [\sigma(\uparrow\uparrow) + \sigma(\downarrow\downarrow)]$ where the arrows denote parallel and antiparallel beam/target spin combinations, either transversely (T) or longitudinally (L) polarized.

Near threshold only a limited number of partial waves are important and considering the allowed transitions to final states⁴ with $L_{NN} \leq 1$ and $l_\pi \leq 1$, i.e. assuming $\sigma_{tot} \equiv \sigma_{Ss} + \sigma_{Ps} + \sigma_{Pp}$, we obtain the following relations⁵

$$^1\sigma_0 = 4\sigma_{Ps} \quad (4)$$

$$^3\sigma_0 = 4(\sigma_{Ss} + \hat{\sigma}_{Pp}) \quad (5)$$

$$^3\sigma_1 = 2(\sigma_{Pp} - \hat{\sigma}_{Pp}) \quad (6)$$

where $\hat{\sigma}_{Pp}$ corresponds to $S = 1$, $m_s = 0$ initial states that contribute to Pp final states but not to σ_{tot} . To which extent the Ps partial waves contribute to σ_{tot} can then be obtained in a model-free way[1] using equ. 1-6

$$\frac{\sigma_{Ps}}{\sigma_{tot}} = \frac{1}{4} \left(1 + \frac{\Delta\sigma_T}{\sigma_{tot}} + \frac{1}{2} \frac{\Delta\sigma_L}{\sigma_{tot}} \right) \quad (7)$$

Theoretical models for pion production near threshold can thus be tested in an unprecedented manner using measurements of spin observables of the reaction $\vec{p}\vec{p} \rightarrow pp\pi^0$ [2]. The results for the spin-correlation coefficient combinations $A_\Delta = A_{xx} - A_{yy}$, and $A_\Sigma = A_{xx} + A_{yy}$ (the latter equivalent to $-\Delta\sigma_T/\sigma_{tot}$) and the analyzing power, A_y , which only needed transverse beam and target polarization, were published recently[3] and the publication of the the data from the experiment using also longitudinal polarization components is in progress[1]. The simultaneous measurements of spin correlation coefficients in $\vec{p}\vec{p} \rightarrow pn\pi^+$ will further enhance the usefulness of this experiment for resolving the theoretical issues of $NN \rightarrow NN\pi$ reactions[4].

Section 2 contains a brief description of the experimental method which is described in detail elsewhere[5,6]. In section 3 we discuss the preliminary results of the measurements using polarized beam with longitudinal components, and the possible implications.

2. THE EXPERIMENT

The vertically polarized beam was injected into the IUCF Cooler ring at 197 MeV and then accelerated to 325, 350, 375 or 400 MeV. The longitudinal component of the beam polarization was achieved using one superconducting solenoid and the three existing solenoids in the cooling section operated in a non-compensating mode. At the injection energy the polarization at the target was almost entirely longitudinal but since the solenoid fields remained constant, after acceleration only partially longitudinal beam polarization could be achieved. The major components were along and vertical to the beam momentum and one minor component was sideways. The beam spin direction was reversed at each new injection.

The target was polarized atomic hydrogen in a gas storage cell fed by an atomic beam source[7]. The target thickness was of the order of 10^{13} atoms/cm². The spin direction was changed every two seconds, along the $\pm x$, $\pm y$ and $\pm z$ axes, by switching weak guide fields ($\sim 0.3 - 0.6$ mT), supplied by Helmholtz-like coils external to the target chamber. A total of 12 beam/target spin combinations were used. The four-momenta of the two

⁴ L_{NN} is the angular momentum between the two nucleons and l_π is the angular momentum of the pion with respect to the nucleon pair.

⁵ Sp is forbidden by parity and angular momentum conservation




Figure 1. The forward detector set-up with an angular coverage of $5^\circ \leq \theta_{lab} \leq 32^\circ$. The different parts are indicated: S: storage cell, F: front detector used for time-of-flight measurements, E, K: stopping scintillators with eight and four segments, thicknesses 10 and 15 cm respectively, V: veto detector, WC1,2: wire chambers, X: four monitor scintillators for measuring pp elastic scattering at $\theta_{lab} = 45^\circ$ and $\phi = \pm 45^\circ$ and $\pm 135^\circ$.




Figure 2. The unpolarized distribution of the missing mass at 325 MeV beam energy in bins of $1 \text{ MeV}/c^2$. The shaded area indicates the background which has been normalized to the area corresponding to invariant masses below $100 \text{ MeV}/c^2$. The filled circles represent the data after the background has been subtracted, the statistical errors are shown. The resolution (FWHM) of the background subtracted peak is $7.2 \text{ MeV}/c^2$.

outgoing protons were measured in a cylindrically symmetric forward detector stack, see Fig. 1, which allowed for the missing mass, m_x , of the third particle to be calculated. In Fig. 2 the m_x distribution is shown for the data set with longitudinal and transverse beam spin components summed over all spin states at a beam energy of 325 MeV. The shape of the background in the spectrum was well reproduced by data taken with nitrogen gas instead of hydrogen in the target.

Full coverage of phase space was achieved with the exception of the loss of events when protons go undetected into the beampipe, i.e. $30 - 22\%$ of phase space distributed events were lost at 325-400 MeV. In an investigation of systematic errors the spin observables were studied as a function of an artificial θ_{lab} cut-off using an effective-range expansion for the pp final state interaction[8]. The corrections applied for the relative spin-dependent cross sections were less than the statistical errors.

The cross-ratio method[9] which eliminates asymmetries due to detector acceptance and spin-dependent luminosities, and division by the product⁶ of the beam and target

⁶ $\Delta\sigma_{T(L)}/\sigma_{tot} = \frac{2}{PQ} \frac{\sqrt{R}-1}{\sqrt{R}+1}$ where R is the ratio of the yields, $Y_{\uparrow\downarrow}Y_{\downarrow\uparrow}/Y_{\uparrow\uparrow}Y_{\downarrow\downarrow}$ and PQ is the product of the beam and target polarization.

polarization projections, $P_y Q_y$ or $P_z Q_z$, give $\Delta\sigma_T/\sigma_{tot}$ and $\Delta\sigma_L/\sigma_{tot}$ from the spinsorted and backgroundsubtracted yields of the π^0 . The values of $P_y Q_y$ and $P_z Q_z$ are obtained by detecting pp elastic scattering at $\theta_{lab} = 45^\circ$ simultaneously with the pion production and measuring the pp spin correlation coefficients A_Δ and A_{zz} which are large and known with high precision at this angle[10–13].

3. PRELIMINARY RESULTS AND DISCUSSION




Figure 3. The data points are the preliminary results for the initial spin state cross sections. The curves are the predictions by the Jülich meson-exchange model (see text). The divisions on the y-axis are arbitrary.

The analysis of the experiment described here is near completion[1]. Preliminary results using equ. 1-3 are shown in Fig. 3. Also depicted are curves representing predictions by a microscopic model calculation, which is based on direct production and off-shell rescattering diagrams. Contributions from heavy meson-exchange (ω) are added and the Δ -isobar is included explicitly[14,15]. Previous comparison with data yielded qualitative agreement for $-A_\Sigma$ and A_y , but disagreed completely with the data on A_Δ [5]. Fig. 3 indicates that ${}^3\sigma_1$, the initial state cross section contributing only to Pp final states, contains the amplitudes quantitatively best reproduced by theory so far.

In [1] it is shown that the energy dependence assumed for the Ps and Pp partial waves, i.e. the proportionality to η^6 and η^8 respectively, is valid for the measured energy range. The magnitude of the Ps cross section is calculated using equ. 7, it increases from about 18% to 30% for bombarding energies in the range from 325 to 400 MeV. At 310 MeV there are recent reports of a significant Ps contribution to the total cross section[16] and an Sd-Ss interference of a few percent[17], measuring unpolarized angular distributions.

We conclude that it should be of high priority to proceed with the study of the main inelastic channel in NN interactions, both experimentally by means of high statistics experiments that allow the extraction of angular distributions of spin-dependent observables, and theoretically by extending the many existing models concerning the Ss contribution to the total cross section to also include higher partial waves.

Acknowledgements

We thank Dr. C. Hanhart for making the theoretical predictions of the model of the Jülich group available and one of the authors, P.T.E., thanks Dr. C. Hanhart for useful discussions and comments. This work has been supported by the US National Science Foundation under Grants PHY95-14566, PHY96-02872, PHY97-22556, and by the US Department of energy under Grant DOE-FG02-88ER40438.

REFERENCES

1. H.O. Meyer et al., submitted to Phys. Rev. Lett., nucl-ex/9907017
2. H.O. Meyer, Proc. 8th Int. Conf. on the Structure of Baryons, Bonn, 22-26 Sep (1998)
3. H.O. Meyer et al., Phys. Rev. Lett. 81 (1998) 3096
4. Swapan K. Saha, W.W. Daehnick et al., Phys. Lett, in print, nucl-ex/9907016
5. P. Thörngren Engblom et al., Proc. Int. Conf. Mesons and Light Nuclei, Prague-Pruhonice, Aug 31-Sep 4 1998, World Scientific, in print, nucl-ex/9810013
6. T. Rinckel et al., submitted to Nucl. Inst. Meth.
7. T. Wise, A.D. Roberts and W. Haeberli, Nucl. Inst. Meth. A336 (1993) 410
8. H.O. Meyer et al., Nucl. Phys. A 539 (1992) 633
9. R.C. Hanna, Proc. 2nd Int. Symp. Polarization Phenomena, eds. P. Huber and H. Schopper, Birkhaeuser, Basel (1966) 280
10. W. Haeberli et al., Phys. Rev. C55 (1997) 597
11. F. Rathmann et al., Phys. Rev. C58 (1998) 658
12. B. von Przewoski et al., Phys. Rev. C58 (1998) 1897
13. B. Lorentz et al., submitted to Phys. Rev.
14. C. Hanhart et al., Phys. Lett. B 444 (1998) 25,
15. C. Hanhart et al., contribution to these proceedings
16. J. Zlomańczuk et al., contribution to these proceedings
17. J. Zlomańczuk et al., Phys. Lett. B436 (1998) 251,